

**East Fork San Gabriel River, Los Angeles County
CAP Workbook Threats Assessment Summary Tables
2008**

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Assessment of Target Viability
East Fork San Gabriel River, Los Angeles County

Double-click opens entry form

				Indicator Ratings									
				Bold = Current				<i>Italics = Desired</i>					
Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating	
1	Egg	Landscape Context	Flow during incubation period	Baseflow in relation to avg. annual daily flow	< 25% of avg. annual daily flow	26-50% of avg. annual daily flow		> 50% of avg. annual daily flow	unimpaired flows	Very Good		Sep-07	
1	Egg	Landscape Context	Non-native species	Non-native egg predators	present throughout watershed	present in >50% of watershed	present in < 50% of watershed	absent				Oct-98	
1	Egg	Landscape Context	Water temperature	Mean weekly avg. temperature in redds	< 5 C. and > 13 C.	11.1-13 C.	10.1-11 C.	6-10 C.					
1	Egg	Condition	Substrate quality	Avg. percent fines (<0.85mm) in potential spawning areas	> 17% fines	11-17% fines	5-10 % fines	< 5% fines	probably little sedimentation	Good		Sep-07	
1	Egg	Condition	Substrate quality	Embeddedness	> 75% embedded	50-75% embedded	25-49% embedded	< 25% embedded	probably low embeddedness	Good		Mar-07	
2	Fry	Landscape Context	Dispersal	Barriers between redds and rearing habitat	complete barrier	partial barriers common	partial barriers scarce	no barriers	no barriers	Very Good		Sep-07	
2	Fry	Landscape Context	Non-native species	Non-native fry predators	present throughout watershed	present > 50% of watershed	present < 50% of watershed	absent				Aug-07	
2	Fry	Landscape Context	Sediment supply	Turbidity (no. days turbidity is > 25 NTUs)	> 30 days during fry development period	20-30 days	10-19 days	< 10 days	probably low turbidity	Good		Sep-07	
2	Fry	Condition	Habitat complexity/refugia	Amount of functional high velocity refuge habitat with flows < 15 cm/sec (boulders, overhanging banks, etc.)	none; watercourse in rearing habitat is channelized	some	common	abundant	mostly natural conditions	Good		Jan-07	
3	Juvenile	Landscape Context	Dispersal	Barriers between rearing habitat and estuary	present			absent	dams; main stem channelized	Poor		Sep-07	
3	Juvenile	Landscape Context	Flow during rearing period	Pool habitat > 3 feet in depth	pools scarce or absent	low abundance of pools	high abundance of pools	high abundance of pools with multiple "refuge" pools (> 5 ft deep)	natural conditions	Good		Sep-07	
3	Juvenile	Landscape Context	Non-native species	Non-native juvenile predators	present throughout watershed	present > 50% watershed	present < 50% watershed	absent				Oct-07	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
3	Juvenile	Landscape Context	Summer flow	Percent of unimpaired median summer baseflow (based on long-term mean monthly discharge)	< 70%	70-90%	> 90%	100% over all IP-km	mostly unimpaired flows	Good		Sep-07	
3	Juvenile	Landscape Context	Water temperature	Median weekly average temperature (MWAT) in potential rearing habitat	> 21 C.	18-21 C.	< 18 C.	< 17 C.				May-95	
3	Juvenile	Condition	Estuarine inflows	Percentage of unimpaired freshwater inflow to estuary (necessary for maintaining brackish water < 15 ppt salinity)	< 25%	25-49%	50-75%	> 75%				Sep-07	
3	Juvenile	Condition	Estuarine inflows	Persistence of hypoxic or anoxic saline layer (> 15 ppt) in potential rearing habitat areas between May and onset of winter rains	3 months	1 month	1 week	< 3 days				Sep-96	
3	Juvenile	Condition	Food availability	Species richness	< 25 taxa	25-29 taxa	30-40 taxa	> 40 taxa				Sep-07	
3	Juvenile	Condition	Habitat complexity/refugia	Instream refugia	absent			present (boulders, overhanging banks, etc.)		Good		Sep-07	
3	Juvenile	Condition	Riparian corridor species composition and structure	Mean percent native, undisturbed composition and structure in 100-foot riparian buffer	< 25%	25-50%	51-75%	historic conditions	Largely natural riparian corridor	Good		Sep-07	
4	Smolt	Landscape Context	Dispersal	Number of days when depths are < 0.4 ft anywhere in migration corridor during outmigration period (March through June)	> 10 days	6-10 days	1-5 days	0 days				Sep-07	
4	Smolt	Landscape Context	Flow for downstream passage March through June	Maximum potential rate of diversion by pumping during April and May (expressed as percent of estimate unimpaired median flow in April)	> 150%	100-150%	50-99%	< 50%				Sep-07	
4	Smolt	Landscape Context	Passage to ocean	Number of days stream mouth is open with adequate flow during outmigration period (March through June)	< 30 days	30-60 days	60-90 days	> 90 days				Sep-96	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
5 Adult	Landscape Context	Dispersal	Accessibility of suitable spawning areas (based on TRT criteria)	accessible sites are clumped in one location or < 25% of all tributaries are accessible	25-50% of all tributaries are accessible	50-75% of all tributaries are accessible	> 75% of all tributaries are accessible	dams; flood control features	Poor		Sep-07	
5 Adult	Landscape Context	Dispersal	Number of days stream mouth is open with adequate flow during entry period (1 November to 1 June)	< 30 days	30-60 days	60-90 days	> 90 days				Sep-96	
5 Adult	Landscape Context	Flow during spawning period (spawning and upstream/downstream passage)	Percent of net discharge (unimpaired flow minus total diversions) occurring between 1 December to 1 June, in all water years	> 10%	6-10%	3-5%	< 3%				Jun-02	
5 Adult	Landscape Context	Water temperature	Median weekly average temperature in migration corridor	> 17 C.	15-16.9 C.	13-14.9 C.	10-12.9 C.				May-96	
5 Adult	Size	Population size	Mean annual adult spawner abundance		TRT criteria for low extinction risk (by watershed)						May-07	
6 Multiple Life Stages	Landscape Context	Barriers/diversions	Stream crossings/stream mile	> two/mile			< two/mile				Oct-07	
6 Multiple Life Stages	Landscape Context	Channel flow and morphology	Percent of total watercourse length channelized	> 25%	16-25%	5-15%	< 5%	probably < 5-15%	Good		Sep-07	
6 Multiple Life Stages	Landscape Context	Fire regime/vegetation maturity	Percent of watershed affected by high intensity fire within previous 100 yrs	> 25%	10-24%	5-9%	< 5%					
6 Multiple Life Stages	Landscape Context	Floodplain connectivity	Floodplain connectivity	< 50% of response reaches in watershed have inundation of historic floodplains by bankfull flows (connectivity)	50-65% of response reaches in watershed demonstrate floodplain connectivity	66-80% of response reaches in watershed demonstrate floodplain connectivity	> 80% of response reaches in watershed demonstrate connectivity	mostly connected	Very Good		Sep-07	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating	
6	Multiple Life Stages	Landscape Context	Historic vs Current Spawning Habitat	Fraction of historic spawning tributaries currently accessible to spawners	< 15% available	16-50% available	51-90% available	>90% available	dams; flood control features	Poor		Sep-07	
6	Multiple Life Stages	Landscape Context	Hydrology	Dry stream reaches	> 75% dry reaches	26-75% dry reaches	1-25% dry reaches	no dry reaches; perennial surface flows	perennial	Very Good		Sep-07	
6	Multiple Life Stages	Landscape Context	Hydrology	Hydrograph	severely modified			natural	mostly natural	Good		Sep-07	
6	Multiple Life Stages	Landscape Context	Land use	Distribution of public ownership along main stem of watercourse	< 25% of land bordering main stem of drainage is publicly owned	25-50%	51-75%	> 75%				Jun-07	
6	Multiple Life Stages	Landscape Context	Land use	Miles of road per square mile of watershed within 100 meters of watercourse	> 1 mi	0.5-1.0 mi	0.1-0.49 mi	< 0.1 mi				Sep-07	
6	Multiple Life Stages	Landscape Context	Land use	Miles of roads per square mile of watershed	> 3.0 mi	2.6-3.0 mi	1.6-2.5 mi	< 1.6 mi				Jun-02	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in agricultural use	> 30%	20-29%	10-19%	< 10%				Jun-02	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in agriculture within 100 meters of watercourse	> 20%	11-20%	5-10%	< 5%					
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in public ownership	< 25 % public ownership	25-50%	51-75%	> 75%				Jan-06	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in urban/residential use	> 25%	10-25%	5-9%	< 5%	probably > 25%	Very Good		Sep-07	
6	Multiple Life Stages	Landscape Context	Water quality	General index of toxicity based on severity of adverse effects on fish	Acute lethal effects (fish kill)	Sublethal effects (reduced growth, altered behavior, etc.)	Toxins detected but no sublethal effects	No toxins or contaminants detected				Sep-07	
6	Multiple Life Stages	Landscape Context	Water quality	Percent total impervious surfaces as % of watershed area	>40%	21-40%	5-20%	< 5%				Sep-07	
6	Multiple Life Stages	Condition	Estuarine habitat quality	Current lagoon area as percentage of historic total area	< 25%	26-50%	51-75%	> 75%				Sep-07	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
6	Multiple Life Stages	Condition	Estuarine habitat quality	Depth, LWD, and other habitat elements (e.g. eelgrass)	depth < 1 meter; LWD and/or overhanging banks absent		depth > 1 meter; LWD and/or overhanging banks present					Sep-96	
6	Multiple Life Stages	Condition	Riparian corridor quality	Riparian canopy cover	< 25% cover	25-49% cover	50-75% cover	> 75% cover	riparian corridor mostly in good condition	Good		Sep-07	
6	Multiple Life Stages	Condition	Riparian corridor quality	Riparian corridor species composition	< 25% native composition	25-50% native composition	50-75% native composition	> 75% native composition				Sep-07	

Overall Viability Summary
East Fork San Gabriel River, Los Angeles County

Summary of Threats										
East Fork San Gabriel River, Los Angeles County										
Threats Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages			Overall Threat Rank
Project-specific threats		1	2	3	4	5	6	7	8	
1	Dams and surface water diversions	Low	Low	Very High	-	Very High	Very High			Very High
2	Channel and/or estuary maintenance, dredging, and vegetation control (incl. flood control activities)	Low	Low	Low	-	-	Low			Low
3	Groundwater extraction	Low	Low	Low	-	-	Low			Low
4	Recreational facilities and activities (ORV use, campgrounds, etc.)	Low	Low	Low	-	-	Low			Low
5	Non-point pollution from roads	Low	Low	-	-	-	-			Low
6	Conversion of watershed lands to row crop agriculture	-	-	-	-	-	Low			Low
7	Urban development	-	-	-	-	-	Low			Low
8	Agricultural effluents	-	-	-	-	-	-			-
9	Artificial lagoon breaching	-	-	-	-	-	-			-
10	Culverts, crossings, and bridges	-	-	-	-	-	-			-
11	Gas, water, and/or other utility pipelines	-	-	-	-	-	-			-
12	Illegal collecting, poaching, and/or unauthorized angling	-	-	-	-	-	-			-
13	Invasive, non-native plants	-	-	-	-	-	-			-
14	Levees and channelization	-	-	-	-	-	-			-
15	Livestock Farming & Ranching	-	-	-	-	-	-			-
16	Mining & Quarrying	-	-	-	-	-	-			-
Threat Status for Targets and Project		Low	Low	High	-	High	High	-	-	Very High *

Overall Viability Summary
East Fork San Gabriel River, Los Angeles County

Summary of Threats										
East Fork San Gabriel River, Los Angeles County										
Threats Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages			Overall Threat Rank
Project-specific threats		1	2	3	4	5	6	7	8	
17	Non-native species present (incl. hatchery fish)	-	-	-	-	-	-			-
18	Oil & Gas Drilling	-	-	-	-	-	-			-
19	Public ownership in watershed									-
20	Roads in watershed and/or within 300 feet of watercourses	-	-	-	-	-	-			-
21	Urban wastewater effluents (incl. industrial and commercial effluents)	-	-	-	-	-	-			-
22	Wildland fires (incl. debris flows following fires)	-	-	-	-	-	-			-
23										-
24										-
25										-
26										-
27										-
28										-
29										-
30										-
31										-
32										-
Threat Status for Targets and Project		Low	Low	High	-	High	High	-	-	Very High *

Overall Viability Summary
East Fork San Gabriel River, Los Angeles County

Stress Matrix									
East Fork San Gabriel River, Los Angeles County									
Stresses (Altered Key Ecological Attributes) Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages		
		1	2	3	4	5	6	7	8
1	Impaired access to rearing and/or spawning habitat	-	-	-	-	-	Very High	-	-
2	Impaired access to spawning areas	-	-	-	-	Very High	-	-	-
3	Impaired access to estuary	-	-	Very High	-	-	-	-	-
4	Impaired riparian habitat quality	-	-	Medium	-	-	-	-	-
5	Altered riparian habitat quality	-	-	-	-	-	Medium	-	-
6	Impaired substrate quality (sedimentation and embeddedness)	Medium	-	-	-	-	-	-	-
7	Impaired instream habitat complexity/refugia	-	-	Medium	-	-	-	-	-
8	Impaired summer base flows	-	-	Medium	-	-	-	-	-
9	Altered sediment supply	-	Medium	-	-	-	-	-	-
10	Impaired habitat complexity/refugia	-	Medium	-	-	-	-	-	-
11	Impaired flows during rearing period	-	-	Medium	-	-	-	-	-
12	Dispersal barriers between redds and rearing habitat	-	Low	-	-	-	-	-	-
13	Altered land use from natural condition	-	-	-	-	-	Low	-	-
14	Altered hydrograph	-	-	-	-	-	Low	-	-
15	Impaired floodplain connectivity	-	-	-	-	-	Low	-	-
16	Altered base flows during incubation	Low	-	-	-	-	-	-	-

Overall Viability Summary
East Fork San Gabriel River, Los Angeles County

Stress Matrix									
East Fork San Gabriel River, Los Angeles County									
Stresses (Altered Key Ecological Attributes) Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages		
		1	2	3	4	5	6	7	8
17	Impaired water temperature in spawning areas	-	-	-	-	-	-	-	-
18	Impaired access to ocean	-	-	-	-	-	-	-	-
19	Impaired access to stream from ocean (stream mouth closed)	-	-	-	-	-	-	-	-
20	Non-native predators	-	-	-	-	-	-	-	-
21	Impaired water temperatures in migration corridor	-	-	-	-	-	-	-	-
22	Low adult population size	-	-	-	-	-	-	-	-
23	Impaired water quality	-	-	-	-	-	-	-	-
24	Impaired food availability	-	-	-	-	-	-	-	-
25	Altered fire regime/recent fire in watershed	-	-	-	-	-	-	-	-
26	Impaired estuarine inflows	-	-	-	-	-	-	-	-
27	Impaired water temperature	-	-	-	-	-	-	-	-
28	Impaired estuarine habitat quality	-	-	-	-	-	-	-	-
29	Non-native egg predators	-	-	-	-	-	-	-	-
30		-	-	-	-	-	-	-	-
31		-	-	-	-	-	-	-	-
32		-	-	-	-	-	-	-	-

**Overall Viability Summary
East Fork San Gabriel River, Los Angeles County**

Overall Viability Summary East Fork San Gabriel River, Los Angeles County								
Conservation Targets		Landscape Context		Condition		Size		Viability Rank
		Grade	Weight	Grade	Weight	Grade	Weight	
1	Egg	Very Good	1	Good	1	-	1	Very Good
2	Fry	Good	1	Good	1	-	1	Good
3	Juvenile	Poor	1	Good	1	-	1	Fair
4	Smolt	-	1	-	1	-	1	-
5	Adult	Poor	1	-	1	-	1	Poor
6	Multiple Life Stages	Poor	1	Good	1	-	1	Fair
7		-	1	-	1	-	1	-
8		-	1	-	1	-	1	-
Project Biodiversity Health Rank								Fair

Overall Viability Summary

East Fork San Gabriel River, Los Angeles County

Detailed Viability Summary											
East Fork San Gabriel River, Los Angeles County											
Conservation Targets		Key Ecological Attributes				Indicators				Calculated Rank	User Override
		Poor	Fair	Good	Very Good	Poor	Fair	Good	Very Good		
1	Egg										Very Good
	Landscape Context				1				1	Very Good	
	Condition			1				2		Good	
	Size									-	
2	Fry										Good
	Landscape Context			1	1			1	1	Good	
	Condition			1				1		Good	
	Size									-	
3	Juvenile										Fair
	Landscape Context	1		2		1		2		Poor	
	Condition			2				2		Good	
	Size									-	
4	Smolt										-
	Landscape Context									-	
	Condition									-	
	Size									-	
5	Adult										Poor
	Landscape Context	1				1				Poor	
	Condition									-	
	Size									-	
6	Multiple Life Stages										Fair
	Landscape Context	1		1	3	1		2	3	Poor	
	Condition			1				1		Good	
	Size									-	
7											-
	Landscape Context									-	
	Condition									-	
	Size									-	
8											-
	Landscape Context									-	
	Condition									-	
	Size									-	