

**San Gabriel River main stem, Los Angeles County
CAP Workbook Threats Assessment Summary Tables
2008**

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Assessment of Target Viability												
San Gabriel River main stem, Los Angeles County												
				Indicator Ratings								
								Double-click opens entry form				
								Bold = Current Italics = Desired				
Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
1	Egg	Landscape Context	Flow during incubation period	Baseflow in relation to avg. annual daily flow	< 25% of avg. annual daily flow	26-50% of avg. annual daily flow	> 50% of avg. annual daily flow				Sep-07	
1	Egg	Landscape Context	Non-native species	Non-native egg predators	present throughout watershed	present in >50% of watershed	present in < 50% of watershed	absent			Oct-98	
1	Egg	Landscape Context	Water temperature	Mean weekly avg. temperature in redds	< 5 C. and > 13 C.	11.1-13 C.	10.1-11 C.	6-10 C.				
1	Egg	Condition	Substrate quality	Avg. percent fines (<0.85mm) in potential spawning areas	> 17% fines	11-17% fines	5-10 % fines	< 5% fines	no spawning habitat left in main stem; tributaries inaccessible	Poor	Sep-07	
1	Egg	Condition	Substrate quality	Embeddedness	> 75% embedded	50-75% embedded	25-49% embedded	< 25% embedded	poor conditions due to periodic sluicing	Poor	May-95	
2	Fry	Landscape Context	Dispersal	Barriers between redds and rearing habitat	complete barrier	partial barriers common	partial barriers scarce	no barriers			Sep-07	
2	Fry	Landscape Context	Non-native species	Non-native fry predators	present throughout watershed	present > 50% of watershed	present < 50% of watershed	absent			Aug-07	
2	Fry	Landscape Context	Sediment supply	Turbidity (no. days turbidity is > 25 NTUs)	> 30 days during fry development period	20-30 days	10-19 days	< 10 days			Sep-07	
2	Fry	Condition	Habitat complexity/refugia	Amount of functional high velocity refuge habitat with flows < 15 cm/sec (boulders, overhanging banks, etc.)	none; watercourse in rearing habitat is channelized	some	common	abundant	main stem below dam is channelized; tributaries are inaccessible	Poor	Jan-07	
3	Juvenile	Landscape Context	Dispersal	Barriers between rearing habitat and estuary	present			absent	main stem channelized; tributaries inaccessible	Poor	Sep-07	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
3	Juvenile	Landscape Context	Flow during rearing period	Pool habitat > 3 feet in depth	pools scarce or absent	low abundance of pools	high abundance of pools	high abundance of pools with multiple "refuge" pools (> 5 ft deep)				Sep-07	
3	Juvenile	Landscape Context	Non-native species	Non-native juvenile predators	present throughout watershed	present > 50% watershed	present < 50% watershed	absent				Oct-07	
3	Juvenile	Landscape Context	Summer flow	Percent of unimpaired median summer baseflow (based on long-term mean monthly discharge)	< 70%	70-90%	> 90%	100% over all IP-km	summer flows depend on surface runoff from roads	Poor		Sep-07	
3	Juvenile	Landscape Context	Water temperature	Median weekly average temperature (MWAT) in potential rearing habitat	> 21 C.	18-21 C.	< 18 C.	< 17 C.				May-95	
3	Juvenile	Condition	Estuarine inflows	Percentage of unimpaired freshwater inflow to estuary (necessary for maintaining brackish water < 15 ppt salinity)	< 25%	25-49%	50-75%	> 75%				Sep-07	
3	Juvenile	Condition	Estuarine inflows	Persistence of hypoxic or anoxic saline layer (> 15 ppt) in potential rearing habitat areas between May and onset of winter rains	3 months	1 month	1 week	< 3 days				Sep-96	
3	Juvenile	Condition	Food availability	Species richness	< 25 taxa	25-29 taxa	30-40 taxa	> 40 taxa				Sep-07	
3	Juvenile	Condition	Habitat complexity/refugia	Instream refugia	absent			present (boulders, overhanging banks, etc.)	main stem channelized, dams; sediment sluicing	Poor		Sep-07	
3	Juvenile	Condition	Riparian corridor species composition and structure	Mean percent native, undisturbed composition and structure in 100-foot riparian buffer	< 25%	25-50%	51-75%	historic conditions	little or no riparian corridor remaining	Poor		Sep-07	
4	Smolt	Landscape Context	Dispersal	Number of days when depths are < 0.4 ft anywhere in migration corridor during outmigration period (March through June)	> 10 days	6-10 days	1-5 days	0 days	highly modified streamflows	Poor		Sep-07	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
4	Smolt	Landscape Context	Flow for downstream passage March through June	Maximum potential rate of diversion by pumping during April and May (expressed as percent of estimate unimpaired median flow in April)	> 150%	100-150%	50-99%	< 50%				Sep-07	
4	Smolt	Landscape Context	Passage to ocean	Number of days stream mouth is open with adequate flow during outmigration period (March through June)	< 30 days	30-60 days	60-90 days	> 90 days				Sep-96	
5	Adult	Landscape Context	Dispersal	Accessibility of suitable spawning areas (based on TRT criteria)	accessible sites are clumped in one location or < 25% of all tributaries are accessible	25-50% of all tributaries are accessible	50-75% of all tributaries are accessible	> 75% of all tributaries are accessible	tributaries inaccessible; main stem channelized	Poor		Sep-07	
5	Adult	Landscape Context	Dispersal	Number of days stream mouth is open with adequate flow during entry period (1 November to 1 June)	< 30 days	30-60 days	60-90 days	> 90 days				Sep-96	
5	Adult	Landscape Context	Flow during spawning period (spawning and upstream/downstream passage)	Percent of net discharge (unimpaired flow minus total diversions) occurring between 1 December to 1 June, in all water years	> 10%	6-10%	3-5%	< 3%				Jun-02	
5	Adult	Landscape Context	Water temperature	Median weekly average temperature in migration corridor	> 17 C.	15-16.9 C.	13-14.9 C.	10-12.9 C.				May-96	
5	Adult	Size	Population size	Mean annual adult spawner abundance		TRT criteria for low extinction risk (by watershed)						May-07	
6	Multiple Life Stages	Landscape Context	Barriers/diversions	Stream crossings/stream mile	> two/mile			< two/mile	avg 2.05 crossings/mile	Poor		Jan-08	
6	Multiple Life Stages	Landscape Context	Channel flow and morphology	Percent of total watercourse length channelized	> 25%	16-25%	5-15%	< 5%	over 52% of main stem	Poor		Sep-07	
6	Multiple Life Stages	Landscape Context	Fire regime/vegetation maturity	Percent of watershed affected by high intensity fire within previous 100 yrs	> 25%	10-24%	5-9%	< 5%	21%	Fair		Jan-08	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating	
6	Multiple Life Stages	Landscape Context	Floodplain connectivity	Floodplain connectivity	< 50% of response reaches in watershed have inundation of historic floodplains by bankfull flows (connectivity)	50-65% of response reaches in watershed demonstrate floodplain connectivity	66-80% of response reaches in watershed demonstrate floodplain connectivity	> 80% of response reaches in watershed demonstrate connectivity	< 48% connectivity	Poor		Sep-07	
6	Multiple Life Stages	Landscape Context	Historic vs Current Spawning Habitat	Fraction of historic spawning tributaries currently accessible to spawners	< 15% available	16-50% available	51-90% available	>90% available	21%; Santa Fe Dam blocks all tributaries	Poor		Sep-07	
6	Multiple Life Stages	Landscape Context	Hydrology	Dry stream reaches	> 75% dry reaches	26-75% dry reaches	1-25% dry reaches	no dry reaches; perennial surface flows				Sep-07	
6	Multiple Life Stages	Landscape Context	Hydrology	Hydrograph	severely modified			natural	highly urbanized watershed	Poor		Sep-07	
6	Multiple Life Stages	Landscape Context	Land use	Distribution of public ownership along main stem of watercourse	< 25% of land bordering main stem of drainage is publicly owned	25-50%	51-75%	> 75%	35%	Fair		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Miles of road per square mile of watershed within 100 meters of watercourse	> 1 mi	0.5-1.0 mi	0.1-0.49 mi	< 0.1 mi	avg 0.98 mi/sq mile	Fair		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Miles of roads per square mile of watershed	> 3.0 mi	2.6-3.0 mi	1.6-2.5 mi	< 1.6 mi	avg 8.9 mi/sq mile	Poor		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in agricultural use	> 30%	20-29%	10-19%	< 10%	1.4%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in agriculture within 100 meters of watercourse	> 20%	11-20%	5-10%	< 5%	0.1%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in public ownership	< 25 % public ownership	25-50%	51-75%	> 75%	35%	Fair		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in urban/residential use	> 25%	10-25%	5-9%	< 5%	53%	Poor		Jan-08	

Assessment of Target Viability
Goleta Slough (San Jose, Atascadero, and Maria Ygnacio creeks, Santa Barbara County)

Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating	
6	Multiple Life Stages	Landscape Context	Water quality	General index of toxicity based on severity of adverse effects on fish	Acute lethal effects (fish kill)	Sublethal effects (reduced growth, altered behavior, etc.)	Toxins detected but no sublethal effects	No toxins or contaminants detected		Fair		Sep-07	
6	Multiple Life Stages	Landscape Context	Water quality	Percent total impervious surfaces as % of watershed area	>40%	21-40%	5-20%	< 5%	27%	Poor		Jan-08	
6	Multiple Life Stages	Condition	Estuarine habitat quality	Current lagoon area as percentage of historic total area	< 25%	26-50%	51-75%	> 75%	2%; lagoon/estuary almost eliminated	Poor		Jan-08	
6	Multiple Life Stages	Condition	Estuarine habitat quality	Depth, LWD, and other habitat elements (e.g. eelgrass)	depth < 1 meter; LWD and/or overhanging banks absent		depth > 1 meter; LWD and/or overhanging banks present					Sep-96	
6	Multiple Life Stages	Condition	Riparian corridor quality	Riparian canopy cover	< 25% cover	25-49% cover	50-75% cover	> 75% cover	35%; most of the main stem on coastal plain is unvegetated	Fair		Jan-08	
6	Multiple Life Stages	Condition	Riparian corridor quality	Riparian corridor species composition	< 25% native composition	25-50% native composition	50-75% native composition	> 75% native composition				Sep-07	

Overall Viability Summary
San Gabriel River main stem, Los Angeles County

Summary of Threats Click the page-down icon ▼ to the right to view more summary tables.										
San Gabriel River main stem, Los Angeles County										
Threats Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages			Overall Threat Rank
		1	2	3	4	5	6	7	8	
1	Channel and/or estuary maintenance, dredging, and vegetation control (incl. flood control activities)	Very High			Very High					
2	Dams and surface water diversions	Very High			Very High					
3	Levees and channelization	Very High			Very High					
4	Urban development	Very High			Very High					
5	Culverts, crossings, and bridges	-	Very High			Very High				
6	Groundwater extraction	-	Very High			Very High				
7	Mining & Quarrying	High	-	Very High	-	-	Very High			Very High
8	Non-point pollution from roads	Very High	-	-	-	-	Very High			Very High
9	Agricultural effluents	-	-	-	-	-	-			-
10	Artificial lagoon breaching	-	-	-	-	-	-			-
11	Conversion of watershed lands to row crop agriculture	-	-	-	-	-	-			-
12	Gas, water, and/or other utility pipelines	-	-	-	-	-	-			-
13	Illegal collecting, poaching, and/or unauthorized angling	-	-	-	-	-	-			-
14	Invasive, non-native plants	-	-	-	-	-	-			-
15	Livestock Farming & Ranching	-	-	-	-	-	-			-
16	Non-native species present (incl. hatchery fish)	-	-	-	-	-	-			-
Threat Status for Targets and Project		Very High	-	-	Very High					

Overall Viability Summary
San Gabriel River main stem, Los Angeles County

Threats Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages			Overall Threat Rank
		1	2	3	4	5	6	7	8	
Project-specific threats		1	2	3	4	5	6	7	8	
17	Oil & Gas Drilling	-	-	-	-	-	-			-
18	Public ownership in watershed									-
19	Recreational facilities and activities (ORV use, campgrounds, etc.)	-	-	-	-	-	-			-
20	Roads in watershed and/or within 300 feet of watercourses	-	-	-	-	-	-			-
21	Urban wastewater effluents (incl. industrial and commercial effluents)	-	-	-	-	-	-			-
22	Wildland fires (incl. debris flows following fires)	-	-	-	-	-	-			-
23										-
24										-
25										-
26										-
27										-
28										-
29										-
30										-
31										-
32										-
Threat Status for Targets and Project		Very High	-	-	Very High					

Overall Viability Summary
San Gabriel River main stem, Los Angeles County

Stress Matrix									
San Gabriel River main stem, Los Angeles County									
Stresses (Altered Key Ecological Attributes) Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages		
		1	2	3	4	5	6	7	8
1	Impaired riparian habitat quality	-	-	Very High	-	-	-	-	-
2	Impaired water quality	-	-	-	-	-	Very High	-	-
3	Altered riparian habitat quality	-	-	-	-	-	Very High	-	-
4	Impaired estuarine habitat quality	-	-	-	-	-	Very High	-	-
5	Altered land use from natural condition	-	-	-	-	-	Very High	-	-
6	Impaired substrate quality (sedimentation and embeddedness)	Very High	-	-	-	-	-	-	-
7	Dispersal barriers between redds and rearing habitat	-	Very High	-	-	-	-	-	-
8	Altered hydrograph	-	-	-	-	-	Very High	-	-
9	Impaired floodplain connectivity	-	-	-	-	-	Very High	-	-
10	Impaired access to rearing and/or spawning habitat	-	-	-	-	-	Very High	-	-
11	Impaired access to estuary	-	-	Very High	-	-	-	-	-
12	Impaired access to spawning areas	-	-	-	-	Very High	-	-	-
13	Impaired summer base flows	-	-	Very High	-	-	-	-	-
14	Impaired access to ocean	-	-	-	Very High	-	-	-	-
15	Impaired instream habitat complexity/refugia	-	-	Very High	-	-	-	-	-
16	Altered fire regime/recent fire in watershed	-	-	-	-	-	High	-	-

Overall Viability Summary
San Gabriel River main stem, Los Angeles County

Stress Matrix									
San Gabriel River main stem, Los Angeles County									
Stresses (Altered Key Ecological Attributes) Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages		
		1	2	3	4	5	6	7	8
17	Altered base flows during incubation	-	-	-	-	-	-	-	-
18	Impaired estuarine inflows	-	-	-	-	-	-	-	-
19	Impaired access to stream from ocean (stream mouth closed)	-	-	-	-	-	-	-	-
20	Impaired water temperature	-	-	-	-	-	-	-	-
21	Impaired water temperatures in migration corridor	-	-	-	-	-	-	-	-
22	Low adult population size	-	-	-	-	-	-	-	-
23	Impaired flows during rearing period	-	-	-	-	-	-	-	-
24	Impaired habitat complexity/refugia	-	-	-	-	-	-	-	-
25	Impaired food availability	-	-	-	-	-	-	-	-
26	Altered sediment supply	-	-	-	-	-	-	-	-
27	Non-native predators	-	-	-	-	-	-	-	-
28	Impaired water temperature in spawning areas	-	-	-	-	-	-	-	-
29	Non-native egg predators	-	-	-	-	-	-	-	-
30		-	-	-	-	-	-	-	-
31		-	-	-	-	-	-	-	-
32		-	-	-	-	-	-	-	-

Overall Viability Summary
San Gabriel River main stem, Los Angeles County

Overall Viability Summary								
San Gabriel River main stem, Los Angeles County								
Conservation Targets		Landscape Context		Condition		Size		Viability Rank
		Grade	Weight	Grade	Weight	Grade	Weight	
1	Egg	-	1	Poor	1	-	1	Poor
2	Fry	-	1	Poor	1	-	1	Poor
3	Juvenile	Poor	1	Poor	1	-	1	Poor
4	Smolt	Poor	1	-	1	-	1	Poor
5	Adult	Poor	1	-	1	-	1	Poor
6	Multiple Life Stages	Poor	1	Poor	1	-	1	Poor
7		-	1	-	1	-	1	-
8		-	1	-	1	-	1	-
Project Biodiversity Health Rank								Poor

Overall Viability Summary
San Gabriel River main stem, Los Angeles County

San Gabriel River main stem, Los Angeles County												
Conservation Targets		Key Ecological Attributes				Indicators				Calculated Rank	User Override	
		Poor	Fair	Good	Very Good	Poor	Fair	Good	Very Good			
1	Egg										Poor	
	Landscape Context										-	
	Condition	1				2					Poor	
	Size										-	
2	Fry										Poor	
	Landscape Context										-	
	Condition	1				1					Poor	
	Size										-	
3	Juvenile										Poor	
	Landscape Context	2				2					Poor	
	Condition	2				2					Poor	
	Size										-	
4	Smolt										Poor	
	Landscape Context	1				1					Poor	
	Condition										-	
	Size										-	
5	Adult										Poor	
	Landscape Context	1				1					Poor	
	Condition										-	
	Size										-	
6	Multiple Life Stages										Poor	
	Landscape Context	5	3			8	5		2		Poor	
	Condition	1	1			1	1				Poor	
	Size										-	
7												-
	Landscape Context										-	
	Condition										-	
	Size										-	
8												-
	Landscape Context										-	
	Condition										-	
	Size										-	