

**San Jose Creek, Monterey County
CAP Workbook Threats Assessment Summary Tables
2008**

Assessment of Target Viability

Double-click opens entry form				Indicator Ratings	Indicator Ratings		Indicator Ratings						
				Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings	Indicator Ratings
Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating	
1 Egg	Landscape Context	Flow during incubation period	Baseflow in relation to avg. annual daily flow	< 25% of avg. annual daily flow	26-50% of avg. annual daily flow		> 50% of avg. annual daily flow	fair; water diversions	Fair		Apr-07		
1 Egg	Landscape Context	Non-native species	Non-native egg predators	present throughout watershed	present in >50% of watershed	present in < 50% of watershed	absent				Aug-07		
1 Egg	Landscape Context	Water temperature	Mean weekly avg. temperature in redds	< 5 C. and > 13 C.	11.1-13 C.	10.1-11 C.	6-10 C.				Oct-05		
1 Egg	Condition	Substrate quality	Avg. percent fines (<0.85mm) in potential spawning areas	> 17% fines	11-17% fines	5-10 % fines	< 5% fines	probably high	Fair		Apr-07		
1 Egg	Condition	Substrate quality	Embeddedness	> 75% embedded	50-75% embedded	25-49% embedded	< 25% embedded	probably high embeddedness	Fair		Mar-03		
2 Fry	Landscape Context	Dispersal	Barriers between redds and rearing habitat	complete barrier	partial barriers common	partial barriers scarce	no barriers	low flows	Fair		Jul-06		
2 Fry	Landscape Context	Non-native species	Non-native fry predators	present throughout watershed	present > 50% watershed	present < 50% of watershed	absent				Aug-07		
2 Fry	Landscape Context	Sediment supply	Turbidity (no. days turbidity is > 25 NTUs)	> 30 days during fry development period	20-30 days	10-19 days	< 10 days	probably high	Fair		Jul-06		
2 Fry	Condition	Habitat complexity/refugia	Amount of functional high velocity refuge habitat with flows < 15 cm/sec (boulders, overhanging banks, etc.)	none; watercourse in rearing habitat is channelized	some	common	abundant				Jul-06		
3 Juvenile	Landscape Context	Dispersal	Barriers between rearing habitat and estuary	present			absent	partial	Fair		Apr-07		
3 Juvenile	Landscape Context	Flow during rearing period	Pool habitat > 3 feet in depth	pools scarce or absent	low abundance of pools	high abundance of pools	high abundance of pools with multiple "refuge" pools (> 5 ft deep)	significant groundwater pumping	Poor		Apr-07		
3 Juvenile	Landscape Context	Non-native species	Non-native juvenile predators	present throughout watershed	present > 50% watershed	present < 50% watershed	absent				Aug-07		
3 Juvenile	Landscape Context	Summer flow	Percent of unimpaired median summer baseflow (based on long-term mean monthly discharge)	< 70%<i>s</i>	70-90%	> 90%	100% over all IP-km	seasonal surface flows	Poor		Apr-07		

Assessment of Target Viability

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
3	Juvenile	Landscape Context	Water temperature	Median weekly average temperature (MWAT) in potential rearing habitat	> 21 C.	18-21 C.	< 18 C.	< 17 C.				Oct-05	
3	Juvenile	Condition	Estuarine inflows	Percentage of unimpaired freshwater inflow to estuary (necessary for maintaining brackish water < 15 ppt salinity)	< 25%	25-49%	50-75%	> 75%	water diversions limit inflows	Poor		Apr-07	
3	Juvenile	Condition	Estuarine inflows	Persistence of hypoxic or anoxic saline layer (> 15 ppt) in potential rearing habitat areas between May and onset of winter rains	3 months	1 month	1 week	< 3 days				Jul-06	
3	Juvenile	Condition	Food availability	Species richness	< 25 taxa	25-29 taxa	30-40 taxa	> 40 taxa				Mar-07	
3	Juvenile	Condition	Habitat complexity/refugia	Instream refugia	absent			present (boulders, overhanging banks, etc.)	limited by seasonal surface flows	Fair		Apr-07	
3	Juvenile	Condition	Riparian corridor species composition and structure	Mean percent native, undisturbed composition and structure in 100-foot riparian buffer	< 25%	25-50%	51-75%	historic conditions				Jul-06	
4	Smolt	Landscape Context	Dispersal	Number of days when depths are < 0.4 ft anywhere in migration corridor during outmigration period (March through June)	> 10 days	6-10 days	1-5 days	0 days	seasonal surface flows	Fair		Apr-07	
4	Smolt	Landscape Context	Flow for downstream passage March through June	Maximum potential rate of diversion by pumping during April and May (expressed as percent of estimate unimpaired median flow in April)	> 150%	100-150%	50-99%	< 50%	significant water diversions	Fair		Apr-07	
4	Smolt	Landscape Context	Passage to ocean	Number of days stream mouth is open with adequate flow during outmigration period (March through June)	< 30 days	30-60 days	60-90 days	> 90 days				Jul-06	
5	Adult	Landscape Context	Dispersal	Accessibility of suitable spawning areas (based on TRT criteria)	accessible sites are clumped in one location or < 25% of all tributaries are accessible	25-50% of all tributaries are accessible	50-75% of all tributaries are accessible	> 75% of all tributaries are accessible	fair	Fair		Apr-07	

Assessment of Target Viability

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
5	Adult	Landscape Context	Dispersal	Number of days stream mouth is open with adequate flow during entry period (1 November to 1 June)	< 30 days	30-60 days	60-90 days	> 90 days				Jul-06	
5	Adult	Landscape Context	Flow during spawning period (spawning and upstream/downstream passage)	Percent of net discharge (unimpaired flow minus total diversions) occurring between 1 December to 1 June, in all water years	> 10%	6-10%	3-5%	< 3%	significant water diversions	Fair		Apr-07	
5	Adult	Landscape Context	Water temperature	Median weekly average temperature in migration corridor	> 17 C.	15-16.9 C.	13-14.9 C.	10-12.9 C.				Oct-05	
5	Adult	Size	Population size	Mean annual adult spawner abundance		TRT criteria for low extinction risk (by watershed)						Mar-03	
6	Multiple Life Stages	Landscape Context	Barriers/diversions	Stream crossings/stream mile	> two/mile			< two/mile	avg 0.81 crossings/mile	Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Channel flow and morphology	Percent of total watercourse length channelized	> 25%	16-25%	5-15%	< 5%	< 5%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Fire regime/vegetation maturity	Percent of watershed affected by high intensity fire within previous 100 yrs	> 25%	10-24%	5-9%	< 5%	0%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Floodplain connectivity	Floodplain connectivity	< 50% of response reaches in watershed have inundation of historic floodplains by bankfull flows (connectivity)	50-65% of response reaches in watershed demonstrate floodplain connectivity	66-80% of response reaches in watershed demonstrate floodplain connectivity	> 80% of response reaches in watershed demonstrate connectivity	> 80%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Historic vs Current Spawning Habitat	Fraction of historic spawning tributaries currently accessible to spawners	< 15% available	16-50% available	51-90% available	>90% available	26%	Fair		Jan-08	
6	Multiple Life Stages	Landscape Context	Hydrology	Dry stream reaches	> 75% dry reaches	26-75% dry reaches	1-25% dry reaches	no dry reaches; perennial surface flows				Apr-07	

Assessment of Target Viability

Conservation Target	Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating	
6	Multiple Life Stages	Landscape Context	Hydrology	Hydrograph	severely modified			natural	significant groundwater pumping	Poor		Apr-07	
6	Multiple Life Stages	Landscape Context	Land use	Distribution of public ownership along main stem of watercourse	< 25% of land bordering main stem of drainage is publicly owned	25-50%	51-75%	> 75%				Jun-07	
6	Multiple Life Stages	Landscape Context	Land use	Miles of road per square mile of watershed within 100 meters of watercourse	> 1 mi	0.5-1.0 mi	0.1-0.49 mi	< 0.1 mi	avg 0.75 mi/sq mi.	Fair		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Miles of roads per square mile of watershed	> 3.0 mi	2.6-3.0 mi	1.6-2.5 mi	< 1.6 mi	1.82 mi/sq mile	Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in agricultural use	> 30%	20-29%	10-19%	< 10%	< 0.1%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in agriculture within 100 meters of watercourse	> 20%	11-20%	5-10%	< 5%	0.1-0.2%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in public ownership	< 25 % public ownership	25-50%	51-75%	> 75%	0.1%	Poor		Mar-07	
6	Multiple Life Stages	Landscape Context	Land use	Percent of watershed area in urban/residential use	> 25%	10-25%	5-9%	< 5%	0.2%	Very Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Water quality	General index of toxicity based on severity of adverse effects on fish	Acute lethal effects (fish kill)	Sublethal effects (reduced growth, altered behavior, etc.)	Toxins detected but no sublethal effects	No toxins or contaminants detected	low avg total N; moderate avg total P	Good		Jan-08	
6	Multiple Life Stages	Landscape Context	Water quality	Percent total impervious surfaces as % of watershed area	>40%	21-40%	5-20%	< 5%	0.1%	Very Good		Jan-08	
6	Multiple Life Stages	Condition	Estuarine habitat quality	Current lagoon area as percentage of historic total area	< 25%	26-50%	51-75%	> 75%				Mar-03	
6	Multiple Life Stages	Condition	Estuarine habitat quality	Depth, LWD, and other habitat elements (e.g. eelgrass)	depth < 1 meter; LWD and/or overhanging banks absent		depth > 1 meter; LWD and/or overhanging banks present					Jul-06	
6	Multiple Life Stages	Condition	Riparian corridor quality	Riparian canopy cover	< 25% cover	25-49% cover	50-75% cover	> 75% cover	91%	Very Good		Jan-08	

Assessment of Target Viability

Conservation Target		Category	Key Attribute	Indicator	Poor	Fair	Good	Very Good	Current Indicator Status	Current Rating	Desired Rating	Date of Current Rating	Date for Desired Rating
6	Multiple Life Stages	Condition	Riparian corridor quality	Riparian corridor species composition	< 25% native composition	25-50% native composition	50-75% native composition	> 75% native composition				Mar-03	

Overall Viability Summary San Jose Creek, Monterey County

Threats Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages			Overall Threat Rank
		1	2	3	4	5	6	7	8	
1	Dams and surface water diversions	High	High	Very High	High	High	High			Very High
2	Groundwater extraction	High	High	Very High	High	High	High			Very High
3	Non-point pollution from roads	High	High	Very High	-	High	Medium			Very High
4	Logging	High	High	High			High			High
5	Culverts, crossings, and bridges	-	-	-	High	High	High			High
6	Urban development	High	-	Medium	-	-	High			High
7	Conversion of watershed lands to row crop agriculture	-	-	-	-	-	High			Medium
8	Roads in watershed and/or within 300 feet of watercourses	-	-	-	-	-	High			Medium
9	Recreational facilities and activities (ORV use, campgrounds, etc.)	-	-	-	-	-	Medium			Low
10	Invasive, non-native plants	-	-	Low	-	-				Low
11	Livestock Farming & Ranching	-	-	-	-	-	Low			Low
12	Agricultural effluents	-	-	-	-	-	-			-
13	Artificial lagoon breaching	-	-	-	-	-	-			-
14	Channel and/or estuary maintenance, dredging, and vegetation control (incl. flood control activities)	-	-	-	-	-	-			-
15	Gas, water, and/or other utility pipelines	-	-	-	-	-	-			-
16	Illegal collecting, poaching, and/or unauthorized angling	-	-	-	-	-	-			-
Threat Status for Targets and Project		High	High	Very High	High	High	Very High	-	-	Very High

**Overall Viability Summary
San Jose Creek, Monterey County**

Threats Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages			Overall Threat Rank
		1	2	3	4	5	6	7	8	
Project-specific threats		1	2	3	4	5	6	7	8	
17	Invasive non-native plants						-			-
18	Levees and channelization	-	-	-	-	-	-			-
19	Log jams and other removable barriers									-
20	Mining & Quarrying	-	-	-	-	-	-			-
21	Natural barriers					-	-			-
22	Non-native species present (incl. hatchery fish)	-	-	-	-	-	-			-
23	Oil & Gas Drilling	-	-	-	-	-	-			-
24	Public ownership in watershed									-
25	Urban wastewater effluents (incl. industrial and commercial effluents)	-	-	-	-	-	-			-
26	Wildland fires (incl. debris flows following fires)	-	-	-	-	-	-			-
27										-

**Overall Viability Summary
San Jose Creek, Monterey County**

Stresses (Altered Key Ecological Attributes) Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages		
		1	2	3	4	5	6	7	8
1	Altered hydrograph	-	-	-	-	-	High	-	-
2	Impaired access to rearing and/or spawning habitat	-	-	-	-	-	High	-	-
3	Altered base flows during incubation	High	-	-	-	-	-	-	-
4	Impaired access to spawning areas	-	-	-	-	High	-	-	-
5	Impaired access to ocean	-	-	-	High	-	-	-	-
6	Impaired substrate quality (sedimentation and embeddedness)	High	-	-	-	-	-	-	-
7	Dispersal barriers between redds and rearing habitat	-	High	-	-	-	-	-	-
8	Impaired instream habitat complexity/refugia	-	-	High	-	-	-	-	-
9	Altered sediment supply	-	High	-	-	-	-	-	-
10	Impaired estuarine inflows	-	-	High	-	-	-	-	-
11	Impaired access to estuary	-	-	High	-	-	-	-	-
12	Impaired flows during rearing period	-	-	High	-	-	-	-	-
13	Impaired summer base flows	-	-	High	-	-	-	-	-
14	Altered land use from natural condition	-	-	-	-	-	Medium	-	-
15	Impaired water quality	-	-	-	-	-	Medium	-	-
16	Altered riparian habitat quality	-	-	-	-	-	Low	-	-

Overall Viability Summary
San Jose Creek, Monterey County

Stress Matrix									
San Jose Creek, Monterey County									
Stresses (Altered Key Ecological Attributes) Across Targets		Egg	Fry	Juvenile	Smolt	Adult	Multiple Life Stages		
		1	2	3	4	5	6	7	8
17	Altered fire regime/recent fire in watershed	-	-	-	-	-	Low	-	-
18	Impaired floodplain connectivity	-	-	-	-	-	Low	-	-
19	Impaired riparian habitat quality	-	-	-	-	-	-	-	-
20	Impaired water temperature in spawning areas	-	-	-	-	-	-	-	-
21	Impaired water temperatures in migration corridor	-	-	-	-	-	-	-	-
22	Low adult population size	-	-	-	-	-	-	-	-
23	Non-native egg predators	-	-	-	-	-	-	-	-
24	Impaired access to stream from ocean (stream mouth closed)	-	-	-	-	-	-	-	-
25	Non-native predators	-	-	-	-	-	-	-	-
26	Impaired food availability	-	-	-	-	-	-	-	-
27	Impaired water temperature	-	-	-	-	-	-	-	-
28	Impaired estuarine habitat quality	-	-	-	-	-	-	-	-
29	Impaired habitat complexity/refugia	-	-	-	-	-	-	-	-
30		-	-	-	-	-	-	-	-
31		-	-	-	-	-	-	-	-
32		-	-	-	-	-	-	-	-

**Overall Viability Summary
San Jose Creek, Monterey County**

Conservation Targets		Landscape Context		Condition		Size		Viability Rank
		Grade	Weight	Grade	Weight	Grade	Weight	
1	Egg	Fair	1	Fair	1	-	1	Fair
2	Fry	Fair	1	-	1	-	1	Fair
3	Juvenile	Poor	1	Poor	1	-	1	Poor
4	Smolt	Fair	1	-	1	-	1	Fair
5	Adult	Fair	1	-	1	-	1	Fair
6	Multiple Life Stages	Poor	1	Very Good	1	-	1	Fair
7		-	1	-	1	-	1	-
8		-	1	-	1	-	1	-
Project Biodiversity Health Rank								Fair

Overall Viability Summary San Jose Creek, Monterey County

Conservation Targets		Key Ecological Attributes				Indicators				Calculated Rank	User Override
		Poor	Fair	Good	Very Good	Poor	Fair	Good	Very Good		
1	Egg	Fair									
	Landscape Context		1				1			Fair	
	Condition		1				2			Fair	
	Size									-	
2	Fry	Fair									
	Landscape Context		2				2			Fair	
	Condition									-	
	Size									-	
3	Juvenile	Poor									
	Landscape Context	2	1			2	1			Poor	
	Condition	1	1			1	1			Poor	
	Size									-	
4	Smolt	Fair									
	Landscape Context		2				2			Fair	
	Condition									-	
	Size									-	
5	Adult	Fair									
	Landscape Context		2				2			Fair	
	Condition									-	
	Size									-	
6	Multiple Life Stages	Fair									
	Landscape Context	1	1	2	4	2	2	3	7	Poor	
	Condition				1				1	Very Good	
	Size									-	
7											
	Landscape Context									-	
	Condition									-	
	Size									-	
8											
	Landscape Context									-	
	Condition									-	
	Size									-	